| | Incidence | | Mortal | Mortality | | |
|----------------------------------|-----------------------|-----------|--------|------------------|--|--|
| | New Cases Annual Rate | | Deaths | Annual Rate | | |
| Year of Diagnosis or Death | Female | Female | Female | Female | | |
| | | ALL RACES | | | | |
| 1988 | 3,360 | 25.2 | 579 | 4.3 | | |
| 1989 | 3,153 | 23.3 | 575 | 4.2 | | |
| 1990 | 3,275 | 23.8 | 566 | 4.1 | | |
| 1991 | 3,309 | 23.8 | 557 | 3.9 | | |
| 1992 | 3,342 | 23.7 | 601 | 4.2 | | |
| 1993 | 3,378 | 23.5 | 585 | 4.0 | | |
| 1994 | 3,406 | 23.5 | 623 | 4.2 | | |
| 1995 | 3,460 | 23.7 | 600 | 4.0 | | |
| 1996 | 3,456 | 23.4 | 585 | 3.8 | | |
| 1997 | 3,588 | 23.7 | 571 | 3.7 | | |
| 1998 | 3,610 | 23.3 | 597 | 3.8 | | |
| 1999 | 3,578 | 22.5 | 655 | 4.0 | | |
| 2000 | 3,603 | 22.1 | 641 | 3.9 | | |
| 2001 | 3,830 | 23.0 | 637 | 3.8 | | |
| 2002 | 3,684 | 21.6 | 698 | 4.0 | | |
| 2003 | 3,653 | 20.9 | 711 | 4.1 | | |
| 2004 | 3,774 | 21.2 | 683 | 3.8 | | |
| 2005 | 3,956 | 21.8 | 713 | 3.9 | | |
| 2006 | 4,077 | 22.0 | 723 | 3.9 | | |
| 2007 | 4,286 | 22.6 | 786 | 4.2 | | |
| 2008 | 4,441 | 22.9 | 853 | 4.4 | | |
| 2009 | 4,716 | 23.6 | 789 | 4.0 | | |
| 2010 | 4,901 | 24.0 | 889 | 4.4 | | |
| 2011 | 5,071 | 24.3 | 910 | 4.3 | | |
| 2012 | 5,389 | 25.2 | 968 | 4.5 | | |
| 2013 | 5,305 | 24.2 | 992 | 4.5 | | |
| 2014 | 5,717 | 25.6 | 1,015 | 4.5 | | |
| 2015 | 5,956 | 26.1 | 1,123 | 4.8 | | |
| 2016 | 6,222 | 26.7 | 1,249 | 5.3 | | |
| 2017 | 6,336 | 26.7 | 1,204 | 5.0 | | |
| | | | (7 | Table Continues) | | |

| | Incidence | | Morta | Mortality | |
|----------------------------------|-----------------------|------------------|--------|------------------|--|
| | New Cases Annual Rate | | Deaths | Annual Rate | |
| Year of Diagnosis or Death | Female | Female | Female | Female | |
| | AS | SIAN/PACIFIC ISI | ANDER | | |
| 1988 | 139 | 14.4 | 20 | 2.4 | |
| 1989 | 131 | 13.3 | 23 | 2.4 | |
| 1990 | 136 | 12.3 | 21 | 2.2 | |
| 1991 | 165 | 14.2 | 25 | 2.3 | |
| 1992 | 185 | 14.2 | 34 | 2.8 | |
| 1993 | 193 | 14.1 | 24 | 1.9 | |
| 1994 | 194 | 13.4 | 27 | 2.1 | |
| 1995 | 216 | 13.9 | 27 | 2.0 | |
| 1996 | 215 | 14.1 | 35 | 2.6 | |
| 1997 | 238 | 14.1 | 39 | 2.7 | |
| 1998 | 286 | 16.6 | 34 | 2.1 | |
| 1999 | 288 | 15.6 | 36 | 2.2 | |
| 2000 | 287 | 14.5 | 39 | 2.2 | |
| 2001 | 348 | 16.7 | 43 | 2.1 | |
| 2002 | 356 | 16.1 | 48 | 2.3 | |
| 2003 | 380 | 16.5 | 65 | 3.1 | |
| 2004 | 421 | 17.2 | 57 | 2.4 | |
| 2005 | 398 | 15.7 | 63 | 2.6 | |
| 2006 | 450 | 17.1 | 64 | 2.5 | |
| 2007 | 482 | 17.5 | 82 | 3.2 | |
| 2008 | 530 | 18.3 | 85 | 3.1 | |
| 2009 | 587 | 19.7 | 78 | 2.7 | |
| 2010 | 610 | 19.6 | 94 | 3.1 | |
| 2011 | 633 | 19.4 | 99 | 3.1 | |
| 2012 | 718 | 21.3 | 118 | 3.6 | |
| 2013 | 730 | 21.2 | 109 | 3.3 | |
| 2014 | 784 | 21.6 | 117 | 3.2 | |
| 2015 | 798 | 21.5 | 136 | 3.6 | |
| 2016 | 835 | 21.4 | 146 | 3.7 | |
| 2017 | 935 | 23.4 | 169 | 4.2 | |
| | | | (| Table Continues) | |

| | Incidence | | Mortality | |
|----------------------------------|-----------|----------------|-----------|------------------|
| | New Cases | Annual Rate | Deaths | Annual Rate |
| Year of Diagnosis or Death | Female | Female | Female | Female |
| | 1 | NON-HISPANIC B | LACK | |
| 1988 | 126 | 16.8 | 42 | 5.8 |
| 1989 | 139 | 18.5 | 54 | 7.3 |
| 1990 | 124 | 15.7 | 49 | 6.8 |
| 1991 | 123 | 15.7 | 49 | 6.2 |
| 1992 | 131 | 16.4 | 47 | 6.1 |
| 1993 | 147 | 17.4 | 61 | 7.7 |
| 1994 | 168 | 19.8 | 56 | 7.0 |
| 1995 | 150 | 17.7 | 41 | 5.1 |
| 1996 | 155 | 17.6 | 56 | 6.8 |
| 1997 | 179 | 20.1 | 50 | 6.0 |
| 1998 | 177 | 19.4 | 54 | 6.2 |
| 1999 | 190 | 20.0 | 52 | 5.7 |
| 2000 | 155 | 16.2 | 66 | 7.2 |
| 2001 | 221 | 22.3 | 57 | 6.0 |
| 2002 | 217 | 21.3 | 72 | 7.4 |
| 2003 | 214 | 20.9 | 84 | 8.6 |
| 2004 | 221 | 20.7 | 75 | 7.6 |
| 2005 | 193 | 18.0 | 65 | 6.2 |
| 2006 | 218 | 19.1 | 79 | 7.5 |
| 2007 | 253 | 22.3 | 82 | 7.4 |
| 2008 | 270 | 23.3 | 78 | 7.0 |
| 2009 | 304 | 26.1 | 91 | 8.1 |
| 2010 | 303 | 24.5 | 104 | 9.1 |
| 2011 | 325 | 25.8 | 102 | 8.3 |
| 2012 | 354 | 27.3 | 98 | 7.7 |
| 2013 | 332 | 24.9 | 105 | 8.2 |
| 2014 | 374 | 27.1 | 116 | 8.7 |
| 2015 | 371 | 26.4 | 125 | 9.3 |
| 2016 | 413 | 29.2 | 140 | 9.9 |
| 2017 | 399 | 27.3 | 125 | 8.9 |
| | | | (| Table Continues) |

| | Incidence | | Morta | Mortality | | |
|----------------------------------|-----------------------|----------|--------|------------------|--|--|
| | New Cases Annual Rate | | Deaths | Annual Rate | | |
| Year of Diagnosis or Death | Female | Female | Female | Female | | |
| | | HISPANIC | | | | |
| 1988 | 276 | 15.2 | 45 | 3.1 | | |
| 1989 | 298 | 17.0 | 49 | 3.1 | | |
| 1990 | 305 | 15.8 | 54 | 3.2 | | |
| 1991 | 334 | 17.2 | 67 | 4.0 | | |
| 1992 | 336 | 16.5 | 69 | 3.7 | | |
| 1993 | 353 | 16.9 | 59 | 3.3 | | |
| 1994 | 338 | 15.3 | 78 | 4.1 | | |
| 1995 | 373 | 16.0 | 75 | 3.7 | | |
| 1996 | 401 | 16.6 | 59 | 2.7 | | |
| 1997 | 403 | 15.9 | 64 | 2.7 | | |
| 1998 | 426 | 16.2 | 80 | 3.7 | | |
| 1999 | 463 | 16.7 | 100 | 4.3 | | |
| 2000 | 527 | 17.7 | 96 | 3.9 | | |
| 2001 | 506 | 16.1 | 106 | 4.1 | | |
| 2002 | 555 | 16.8 | 101 | 3.9 | | |
| 2003 | 582 | 16.8 | 102 | 3.5 | | |
| 2004 | 649 | 17.7 | 99 | 3.1 | | |
| 2005 | 694 | 17.8 | 135 | 4.2 | | |
| 2006 | 732 | 17.9 | 118 | 3.4 | | |
| 2007 | 768 | 18.1 | 126 | 3.5 | | |
| 2008 | 815 | 18.1 | 137 | 3.6 | | |
| 2009 | 890 | 19.2 | 159 | 4.0 | | |
| 2010 | 928 | 19.2 | 165 | 3.8 | | |
| 2011 | 1,086 | 21.4 | 180 | 4.1 | | |
| 2012 | 1,184 | 22.1 | 191 | 4.2 | | |
| 2013 | 1,193 | 21.9 | 202 | 4.2 | | |
| 2014 | 1,262 | 22.0 | 201 | 4.0 | | |
| 2015 | 1,462 | 24.4 | 252 | 4.6 | | |
| 2016 | 1,521 | 24.6 | 267 | 4.7 | | |
| 2017 | 1,596 | 25.0 | 290 | 5.0 | | |
| | | | (| Table Continues) | | |

| | Incidence | | Morta | Mortality | |
|----------------------------------|-----------------------|----------------|--------|-------------|--|
| | New Cases Annual Rate | | Deaths | Annual Rate | |
| Year of Diagnosis or Death | Female | Female | Female | Female | |
| | | NON-HISPANIC V | VHITE | | |
| 1988 | 2,788 | 28.1 | 471 | 4.5 | |
| 1989 | 2,569 | 25.7 | 446 | 4.2 | |
| 1990 | 2,683 | 26.5 | 438 | 4.1 | |
| 1991 | 2,660 | 26.4 | 415 | 3.8 | |
| 1992 | 2,662 | 26.3 | 448 | 4.1 | |
| 1993 | 2,664 | 26.1 | 440 | 4.1 | |
| 1994 | 2,669 | 26.3 | 461 | 4.3 | |
| 1995 | 2,684 | 26.7 | 456 | 4.2 | |
| 1996 | 2,642 | 26.2 | 433 | 3.9 | |
| 1997 | 2,741 | 26.8 | 418 | 3.7 | |
| 1998 | 2,692 | 26.2 | 428 | 3.8 | |
| 1999 | 2,611 | 25.2 | 467 | 4.1 | |
| 2000 | 2,604 | 24.8 | 437 | 3.8 | |
| 2001 | 2,722 | 25.7 | 430 | 3.8 | |
| 2002 | 2,509 | 23.5 | 473 | 4.1 | |
| 2003 | 2,434 | 22.6 | 459 | 4.0 | |
| 2004 | 2,453 | 22.7 | 450 | 3.8 | |
| 2005 | 2,631 | 24.4 | 447 | 3.8 | |
| 2006 | 2,638 | 24.2 | 461 | 4.0 | |
| 2007 | 2,724 | 24.7 | 492 | 4.2 | |
| 2008 | 2,773 | 25.1 | 548 | 4.7 | |
| 2009 | 2,874 | 25.5 | 458 | 3.9 | |
| 2010 | 3,003 | 26.4 | 521 | 4.3 | |
| 2011 | 2,966 | 25.7 | 526 | 4.2 | |
| 2012 | 3,063 | 26.5 | 555 | 4.4 | |
| 2013 | 2,961 | 25.1 | 570 | 4.6 | |
| 2014 | 3,200 | 26.9 | 574 | 4.5 | |
| 2015 | 3,225 | 26.9 | 600 | 4.7 | |
| 2016 | 3,333 | 27.5 | 686 | 5.3 | |
| 2017 | 3,258 | 26.5 | 610 | 4.7 | |

| | Incidence | | Mortality | | |
|---------------------------------|---------------------|--------------------|------------------|-------------------|--|
| | New Cases 2013-2017 | Annual Rate | Deaths 2013-2017 | Annual Rate | |
| Age at Diagnosis or Death | Female | Female | Female | Female | |
| | | ALL RACES | | | |
| 00 - 04 | 0 | 0.0 | 0 | 0.0 | |
| 05 - 09 | 0 | 0.0 | 0 | 0.0 | |
| 10 - 14 | 0 | 0.0 | 0 | 0.0 | |
| 15 - 19 | 7 | 0.1 | 0 | 0.0 | |
| 20 - 24 | 35 | 0.5 | 2 | 0.0 | |
| 25 - 29 | 135 | 1.9 | 10 | 0.1 | |
| 30 - 34 | 385 | 5.6 | 24 | 0.4 | |
| 35 - 39 | 648 | 10.0 | 48 | 0.7 | |
| 40 - 44 | 1,029 | 16.0 | 69 | 1.1 | |
| 45 - 49 | 1,642 | 25.4 | 159 | 2.5 | |
| 50 - 54 | 3,110 | 47.2 | 336 | 5.1 | |
| 55 - 59 | 4,761 | 75.2 | 541 | 8.5 | |
| 60 - 64 | 5,229 | 94.7 | 799 | 14.5 | |
| 65 - 69 | 4,839 | 106.1 | 935 | 20.5 | |
| 70 - 74 | 3,260 | 98.2 | 763 | 23.0 | |
| 75 - 79 | 2,057 | 85.0 | 658 | 27.2 | |
| 80 - 84 | 1,322 | 72.8 | 535 | 29.5 | |
| 85 + | 1,077 | 48.1 | 704 | 31.4 | |
| Total/Crude | 29,536 | 30.1 | 5,583 | 5.7 | |
| | | ASIAN/PACIFIC ISLA | ANDER | | |
| 00 - 04 | 0 | 0.0 | 0 | 0.0 | |
| 05 - 09 | 0 | 0.0 | 0 | 0.0 | |
| 10 - 14 | 0 | 0.0 | 0 | 0.0 | |
| 15 - 19 | 1 | 0.1 | 0 | 0.0 | |
| 20 - 24 | 3 | 0.3 | 0 | 0.0 | |
| 25 - 29 | 21 | 1.7 | 1 | 0.1 | |
| 30 - 34 | 70 | 5.7 | 5 | 0.4 | |
| 35 - 39 | 128 | 10.8 | 8 | 0.7 | |
| 40 - 44 | 225 | 18.5 | 11 | 0.9 | |
| 45 - 49 | 366 | 32.2 | 31 | 2.7 | |
| 50 - 54 | 612 | 55.9 | 67 | 6.1 | |
| 55 - 59 | 729 | 69.6 | 92 | 8.8 | |
| 60 - 64 | 652 | 69.8 | 96 | 10.3 | |
| 65 - 69 | 519 | 68.8 | 102 | 13.5 | |
| 70 - 74 | 339 | 65.2 | 83 | 16.0 | |
| 75 - 79 | 201 | 50.9 | 69 | 17.5 | |
| 80 - 84 | 123 | 42.0 | 43 | 14.7 | |
| 85 + | 93 | 28.9 | 69 | 21.4 | |
| Total/Crude | 4,082 | 26.4 | 677 | 4.4 | |
| | | | | (Table Continues) | |

Five-Year Incidence and Mortality Counts and Average Annual Age-Specific and Crude Rates per 100,000 Females by Age and Race/Ethnicity, California, 2013-2017, Corpus Uteri Cancer and Uterus, NOS (continued)

| | Incidence | | Mortality | | |
|---------------------------------|---------------------|-----------------|------------------|-------------------|--|
| | New Cases 2013-2017 | Annual Rate | Deaths 2013-2017 | Annual Rate | |
| Age at Diagnosis or Death | Female | Female | Female | Female | |
| | | NON-HISPANIC BL | ACK | | |
| 00 - 04 | 0 | 0.0 | 0 | 0.0 | |
| 05 - 09 | 0 | 0.0 | 0 | 0.0 | |
| 10 - 14 | 0 | 0.0 | 0 | 0.0 | |
| 15 - 19 | 0 | 0.0 | 0 | 0.0 | |
| 20 - 24 | 3 | 0.6 | 0 | 0.0 | |
| 25 - 29 | 8 | 1.7 | 1 | 0.2 | |
| 30 - 34 | 20 | 4.9 | 2 | 0.5 | |
| 35 - 39 | 23 | 6.0 | 4 | 1.1 | |
| 40 - 44 | 39 | 10.4 | 7 | 1.9 | |
| 45 - 49 | 75 | 18.4 | 8 | 2.0 | |
| 50 - 54 | 133 | 30.2 | 28 | 6.4 | |
| 55 - 59 | 294 | 70.4 | 64 | 15.3 | |
| 60 - 64 | 394 | 115.9 | 114 | 33.5 | |
| 65 - 69 | 376 | 141.8 | 123 | 46.4 | |
| 70 - 74 | 253 | 134.4 | 101 | 53.6 | |
| 75 - 79 | 134 | 95.3 | 75 | 53.4 | |
| 80 - 84 | 84 | 86.8 | 47 | 48.6 | |
| 85 + | 53 | 50.6 | 37 | 35.3 | |
| Total/Crude | 1,889 | 30.9 | 611 | 10.0 | |
| | | HISPANIC | | | |
| 00 - 04 | 0 | 0.0 | 0 | 0.0 | |
| 05 - 09 | 0 | 0.0 | 0 | 0.0 | |
| 10 - 14 | 0 | 0.0 | 0 | 0.0 | |
| 15 - 19 | 3 | 0.1 | 0 | 0.0 | |
| 20 - 24 | 22 | 0.7 | 1 | 0.0 | |
| 25 - 29 | 75 | 2.5 | 7 | 0.2 | |
| 30 - 34 | 208 | 7.4 | 12 | 0.4 | |
| 35 - 39 | 330 | 12.3 | 27 | 1.0 | |
| 40 - 44 | 469 | 18.1 | 34 | 1.3 | |
| 45 - 49 | 614 | 26.0 | 69 | 2.9 | |
| 50 - 54 | 927 | 44.8 | 104 | 5.0 | |
| 55 - 59 | 1,184 | 68.8 | 138 | 8.0 | |
| 60 - 64 | 1,118 | 84.7 | 197 | 14.9 | |
| 65 - 69 | 889 | 92.6 | 185 | 19.3 | |
| 70 - 74 | 537 | 79.8 | 153 | 22.7 | |
| 75 - 79 | 324 | 67.0 | 96 | 19.9 | |
| 80 - 84 | 203 | 58.5 | 84 | 24.2 | |
| 85 + | 131 | 36.0 | 105 | 28.9 | |
| Total/Crude | 7,034 | 18.7 | 1,212 | 3.2 | |
| | | | | (Table Continues) | |

Five-Year Incidence and Mortality Counts and Average Annual Age-Specific and Crude Rates per 100,000 Females by Age and Race/Ethnicity, California, 2013-2017, Corpus Uteri Cancer and Uterus, NOS (continued)

| | Incidence | | <u>Mortality</u> | | |
|---------------------------------|---------------------|----------------|------------------|-------------|--|
| | New Cases 2013-2017 | Annual Rate | Deaths 2013-2017 | Annual Rate | |
| Age at Diagnosis or Death | Female | Female | Female | Female | |
| | | NON-HISPANIC W | /HITE | | |
| 00 - 04 | 0 | 0.0 | 0 | 0.0 | |
| 05 - 09 | 0 | 0.0 | 0 | 0.0 | |
| 10 - 14 | 0 | 0.0 | 0 | 0.0 | |
| 15 - 19 | 2 | 0.1 | 0 | 0.0 | |
| 20 - 24 | 6 | 0.3 | 1 | 0.1 | |
| 25 - 29 | 25 | 1.0 | 1 | 0.0 | |
| 30 - 34 | 76 | 3.1 | 5 | 0.2 | |
| 35 - 39 | 150 | 6.9 | 8 | 0.4 | |
| 40 - 44 | 271 | 12.3 | 16 | 0.7 | |
| 45 - 49 | 551 | 21.7 | 47 | 1.9 | |
| 50 - 54 | 1,365 | 46.4 | 136 | 4.6 | |
| 55 - 59 | 2,457 | 79.2 | 238 | 7.7 | |
| 60 - 64 | 2,974 | 102.8 | 381 | 13.2 | |
| 65 - 69 | 2,980 | 116.7 | 520 | 20.4 | |
| 70 - 74 | 2,076 | 108.2 | 422 | 22.0 | |
| 75 - 79 | 1,371 | 98.8 | 414 | 29.8 | |
| 80 - 84 | 889 | 82.9 | 360 | 33.6 | |
| 85 + | 784 | 54.4 | 491 | 34.1 | |
| Total/Crude | 15,977 | 41.5 | 3,040 | 7.9 | |

Risk (Percent) of Being Diagnosed with Invasive Cancer within the Next 10, 20, or 30 Years, or in the Remaining Lifetime, Assuming Cancer Free at Current Age, and Lifetime Risk of Being Diagnosed with or Dying of Cancer by Race/Ethnicity, Corpus Uteri Cancer and Uterus, NOS

Females

| | remaies | | | |
|--|--------------|------------------------------|----------------|------------|
| Current Age | 10+ yrs | 20+ yrs | 30+ yrs | eventually |
| | A | LL RACES | | |
| 0 | 0.00 | 0.00 | 0.01 | 2.86 |
| 25 | 0.04 | 0.16 | 0.52 | 2.88 |
| 35 | 0.13 | 0.48 | 1.25 | 2.86 |
| 45 | 0.36 | 1.14 | 1.99 | 2.76 |
| 55 | 0.80 | 1.68 | 2.23 | 2.47 |
| 65 | 0.93 | 1.52 | - | 1.78 |
| | | isk of Being isk of Dying | | |
| | ASIAN/P | ACIFIC ISL | ANDER | |
| 0 | 0.00 | 0.00 | 0.01 | 2.42 |
| 25 | 0.04 | 0.18 | 0.60 | 2.44 |
| 35 | 0.15 | 0.57 | 1.21 | 2.41 |
| 45 | 0.42 | 1.08 | 1.67 | 2.28 |
| 55 | 0.66 | 1.27 | 1.64 | 1.89 |
| 65 | 0.63 | 1.02 | - | 1.27 |
| | | isk of Being isk of Dying | | |
| | | ISPANIC BI | | |
| 0 | 0.00 | 0.00 | 0.01 | 2.78 |
| 25 | 0.03 | 0.11 | 0.35 | 2.84 |
| 35 | 0.08 | 0.32 | 1.12 | 2.83 |
| 45 | 0.25 | 1.06 | 2.10 | 2.81 |
| 55 | 0.86 | 1.94 | 2.49 | 2.69 |
| 65 | 1.21 | 1.82 | - | 2.05 |
| | | isk of Being isk of Dying | | |
| | | HISPANIC | or Caricer | = 1.0576 |
| 0 | | 0.00 | 0.02 | 2.59 |
| 25 | 0.00 0.05 | 0.00 | 0.02 | 2.59 |
| 35 | 0.15 | 0.49 | 1.19 | 2.58 |
| 45 | 0.35 | 1.06 | 1.79 | 2.45 |
| 55 | 0.72 | 1.47 | 1.92 | 2.15 |
| 65 | 0.79 | 1.27 | - | 1.51 |
| 00 | | isk of Being | Diagnosed | |
| | Lifetime R | isk of Dying | of Cancer | |
| | | IISPANIC W | | |
| 0 | 0.00 | 0.00 | 0.01 | 2.97 |
| 25 | 0.02 | 0.11 | 0.44 | 2.99 |
| 35 | 0.09 | 0.43 | 1.25 | 2.99 |
| 45 55 | 0.34 | 1.17 | 2.10 | 2.93 |
| 55 65 | 0.86 | 1.81 | 2.43 | 2.67 |
| 65 | 1.02 | 1.68 | - Diagnosco | 1.94 |
| Lifetime Risk of Being Diagnosed = 2.97% Lifetime Risk of Dying of Cancer = 0.67% | | | | |