

Inyo and Mono Counties: Status Update of Cancer Incidence, Mortality, and Health Behaviors

December 2015

This fact sheet is provided by the California Cancer Registry (CCR) of the California Department of Public Health to show the intersection of cancer and poor health behaviors on California residents. CCR is California's population-based cancer surveillance system, and it collects information on nearly all cancers diagnosed in the state. This information helps to further our understanding of cancer and it is used to develop strategies and policies for its prevention, treatment, and control.

Most Common Cancers and Cancer Related Deaths by Sex, Inyo and Mono Counties, 2008-2012

INCIDENCE					
MEN	County Rate	State Rate	WOMEN	County Rate	State Rate
1. Prostate	143.4	126.9	1. Breast	128.5	122.1
2. Colon and Rectum	40.9	46.0	2. Lung and Bronchus	41.9	42.1
3. Lung and Bronchus	39.1	55.8	3. Colon and Rectum	26.3	35.1
4. Kidney and Renal Pelvis	26.9	19.6	4. Uterus	17.7	23.3
5. Non-Hodgkin Lymphoma	22.5	22.8	5. Melanoma	16.8	15.6
All Sites	416.5	476.7	All Sites	349.0	388.8


MORTALITY					
MEN	County Rate	State Rate	WOMEN	County Rate	State Rate
1. Lung and Bronchus	22.2	43.5	1. Lung and Bronchus	31.3	30.4
2. Prostate	20.6	21.1	2. Breast	17.9	21.2
3. Colon and Rectum	^	16.7	3. Pancreas	^	9.3
4. Pancreas	^	11.7	4. Colon and Rectum	^	12.1
5. Liver and Intrahepatic Bile Duct	^	10.5	5. Esophagus	^	1.4
All Malignant Cancers	121.5	182.7	All Malignant Cancers	101.7	134.8

Rates are shown as the number of new cases or deaths per 100,000 persons. All rates are age-adjusted to the 2000 United States Standard Population.

^ Rates based on fewer than 15 cases have been suppressed.

Confidence intervals can be obtained from the CCR Data and Mapping tool (<http://www.cancer-rates.info/ca/>) and may help to assess statistical significance of age-adjusted rates.

Age-Adjusted Incidence Rates by Race/Ethnicity and Sex, Inyo and Mono Counties, All Cancer Sites Combined, 2008-2012




Source: California Cancer Registry, California Department of Public Health.

^ Rates based on counts less than 15 have been suppressed.

For more information, please visit the CCR website at www.ccrca.org.


Age-Adjusted Mortality Rates by Race/Ethnicity and Sex, Inyo and Mono Counties, All Cancer Sites Combined, 2008-2012



Source: California Cancer Registry, California Department of Public Health.
 ^ Rates based on counts less than 15 have been suppressed.

Poor lifestyle choices such as overuse of alcohol, smoking, poor diet, and lack of physical exercise are significant contributors in the development and progression of preventable chronic diseases, including several types of cancer. Modification of negative risk behaviors and an increased awareness towards maintaining healthy lifestyle behaviors are vital to the prevention and/or early treatment for improvement of chronic disease outcomes. The California Behavioral Risk Factor Surveillance System (BRFSS) is an annual statewide telephone survey developed to collect data on the health-risk behaviors and use of preventative services among adults in California. BRFSS data are used to help target and build health promotion activities around the state.

Prevalence of Health Risk Behaviors* by Sex, California, 2012



Source: 2012 California Behavioral Risk Factor Surveillance System, California Department of Public Health.

* Data are weighted to the 2010 California population. Obesity is based on Body Mass Index (BMI) ≥ 30 . Colorectal cancer (CRC) screened is defined as having been screened by use of either a sigmoidoscopy in the last five years, a colonoscopy in the last ten years, or by a fecal occult blood test (FOBT) home kit in the last year.